



Sexually Transmitted Diseases (STD's)



June, 2005







Outline

- 1. Why worry?
- 2. The STD Infections
 - Chlamydia & Gonorrhea
 - Syphilis

HIV/AIDS

HPV

Herpes

- Others
- 3. How to prevent STD's
- 4. Getting Tested: Who/Where/How?



Why Worry?

- Rates are going <u>UP</u>:
 - Chlamydia
 - Syphilis
 - HIV
- Teens more likely to catch an infection
- Hispanics & blacks have higher rates





Chlamydia & Gonorrhea





- Caused by bacteria
- Spread person-to-person during sexual activity
- Infection can include many parts of the body
 - Cervix, urethra, genital tract
- Throat infection possible from oral sex
- Curable with antibiotics

Chlamydia & Gonorrhea in FEMALES




- Usually NO SYMPTOMS— Most females feel FI NE
- May have:
 - Vaginal discharge (fluid/mucus)
 - Burning or pain with peeing
 - Lower abdominal pain



Chlamydia & Gonorrhea in FEMALES

- Infection in cervix
- Can go into uterus, tubes, ovaries ("PI D")









Chlamydia & Gonorrhea Can Cause:

- Painful Pelvic Inflammatory Disease (PID)
- Trouble getting pregnant later in life
- Dangerous tubal pregnancy



Chlamydia & Gonorrhea in MALES

Usually NO SYMPTOMS

Urethral discharge (drip)

Burning on peeing

Testicular pain



Gonorrhea



Chlamydia & Gonorrhea Can Cause:

- Irritation to urinary tract
- Infection of the tubes and testicles inside the body
- Trouble getting a woman pregnant later in life


Syphilis

- A bacteria spread by sexual contact.
- Stages of disease
 - Primary- painless sore anywhere sex occurred
 - Secondary body rash
- Many have NO symptoms
- Curable with antibiotics





Syphilis: If not Treated



- Infection continues
 - Even though symptoms go away
- Serious problems years later
 - Heart problems
 - Memory loss
 - Stroke-like problems
- Babies (if Mother infected):
 - Stillborn, Birth defects,
 - Mental retardation

It is much easier to catch HIV/AIDS when infected with Syphilis



HIV/AIDS: Know the Facts

- HIV = a virus
- Spread by blood or body fluids during
 - Sex or Sharing Needles
 - Babies infected at birth
- Virus keeps growing in the body and attacks the immune system.
- Usually no symptoms
- After years, AIDS can develop




NIGHT SWEATS



HIV/AIDS: What would I feel?





Most HIV-infected people feel FINE

Some may have "flu" symptoms weeks after getting infected

If AIDS develops, may get symptoms of AIDS

The only way to know for sure if you have HIV is to GET TÉSTED




HIV/AIDS: Living with It

HIV infection can get worse: AIDS

- Body loses ability to fight infection
- · Severe weight loss, diarrhea
- Memory Problems
- Pneumonia, other serious infections
- Death

Treatable, but not curable Many people on medicines live relatively healthy lives





Genital Herpes

- Common: 1 in 5 adults
- Viral infection of skin/membranes, nerves
- Painful blisters, often come back
- Often no symptoms
- Can transmit even when no ulcers
- No cure, but medicines can help



Trichomonas ("Tric")

- One-cell protozoan parasite
- Very common
- · Women:
 - vaginal discharge, inflammation, odor
- Men often no symptoms
- Cured with antibiotics



HPV: Human Papillomavirus

- Many virus types
- Some cause warts
- Some genital warts
- Others can lead to cancer
 - Cervix (uterus)
 - Penis
 - Anus
 - Throat









Many Other STD's

- Hepatitis B*
- Hepatitis C*
- Pubic Lice*
- Chancroid
- LGV
- Bacterial Vaginosis (BV)*
- Scabies*
- Molluscum
 Contagiousum*

*can be spread by other means than sex



- Pain
- Blisters
- Bumps
- Sores
- Drip or discharge
- I tching
- Scarring
- More likely to get other infections




What can I do not to get an STD or HIV???

- 1. ABSTINENCE—do not have sex (vaginal, oral, or anal)
- 2. Condoms EVERY time
- 3. Few partners
- 4. Low-risk partners
- 5. Lifetime relationship with uninfected partner (BOTH faithful)
- 6. GET CHECKED

Who Should be Tested?

- Anyone who has had sex should be tested
 - Including oral, anal, and vaginal sex
- Anyone who has shared needles or used injection drugs







Testing is Confidential


- Teens can be tested for STD's confidentially.
- Parent permission NOT needed, but discussion with parents always encouraged.
- Information is private.











What Happens during an STD Test?

At Project WORTH and Metro Health Clinics

- Answer questions <u>honestly</u> on paper
- Reviewed by nurse or doctor
- Give blood sample for HIV and Syphilis test
- Give urine sample for Chlamydia & Gonorrhea test





What Happens during an STD Test?

Interview and examination, if needed

Get medicine, if needed

Call or come back for test results

Make follow up appointment if needed

- Other places may do other tests:
 - -Finger-stick for HIV
 - -Oral swab for HIV
 - -Physical exam for all patients
 - -Provide Pap smear for women at visit



Where Can I Get Tested?

- SAMHD clinics
 - Call 207-8850
- Other Community Organizations
 - Beat AIDS
 - 227-4689
 - Hope Action Care
 - 224-7330
 - Community Clinic
 - 821-5522
- Your Primary Doctor



Questions & Answers





Top 10 Questions About STD's



- 1. I know I don't have an STD because we were each other's "first", right?
- 2. Can you get an STD from kissing?
- 3. Can you get an STD from toilet seats or hot tubs?



- 4. Can STD's "come back" after treatment?
- 5. If I had an STD before, I can't get pregnant, right?

Top 10 Questions About STD's

- 6. I know everyone my girlfriend has been with and they are "clean". Why would I need a test?
- 7. I have always used a condom, so I don't need a test, right?





- 8. If an STD test I got at a clinic showed I was infected, they would have called me, right?
- 9. When my boyfriend was in juvenile detention, he was tested for "everything", right?
- 10. I hear getting a check up is expensive. How much will it cost?